

Molecular crowding limits translation and cell growth

Stefan Klumpp^{a,1}, Matthew Scott^b, Steen Pedersen^c, and Terence Hwa^{d,e}

^aMax Planck Institute of Colloids and Interfaces, 14424 Potsdam, Germany; ^bDepartment of Applied Mathematics, University of Waterloo, Waterloo, ON, Canada N2L 3G1; ^cDepartment of Biology, University of Copenhagen, 2200 Copenhagen N, Denmark; and ^dDepartment of Physics and Center for Theoretical Biological Physics, and ^eSection of Molecular Biology, Division of Biological Sciences, University of California, San Diego, La Jolla, CA 92093

Edited by David R. Nelson, Harvard University, Cambridge, MA, and approved September 5, 2013 (received for review May 31, 2013)

Bacterial growth is crucially dependent on protein synthesis and thus on the cellular abundance of ribosomes and related proteins. Here, we show that the slow diffusion of the bulky tRNA complexes in the crowded cytoplasm imposes a physical limit on the speed of translation, which ultimately limits the rate of cell growth. To study the required allocation of ancillary translational proteins to alleviate the effect of molecular crowding, we develop a model for cell growth based on a coarse-grained partitioning of the proteome. We find that coregulation of ribosome- and tRNA-affiliated proteins is consistent with measured growth-rate dependencies and results in near-optimal allocation over a broad range of growth rates. The analysis further resolves a long-standing controversy in bacterial growth physiology concerning the growth-rate dependence of translation speed and serves as a caution against premature identification of phenomenological parameters with mechanistic processes.

Bacterial cell growth and protein synthesis are tightly coupled as proteins account for a large fraction of the cellular biomass (1). In the model organism *Escherichia coli*, over half of the biomass is protein (2), and protein synthesis accounts for more than two-thirds of the cell's ATP budget during rapid growth (3). Therefore, the machinery of protein synthesis, i.e., ribosomes, tRNAs, and ribosome-affiliated factors, plays a central role in maintaining exponential growth (1, 4). This is manifested by an increased ribosome content in rapidly growing cells (2, 5, 6), by direct observations that protein synthesis is limited by the availability of free ribosomes (7), and by considerations that link evolutionary selective pressure to the cost of protein synthesis (8).

The most striking evidence for the central role of ribosomes in cell growth is provided by the linear relation between the ribosome mass fraction and the growth rate for bacteria grown in media containing different nutrients. This linear relation, which emerged from the systematic characterization of bacterial cells growing at different rates (5, 9), is illustrated in Fig. 1A with data for *E. coli* (2, 10, 11). It can be interpreted as reflecting the intrinsically autocatalytic activity of ribosomes synthesizing ribosomal proteins (9, 12) and identifies the fraction of ribosomes allocated to making ribosomal proteins as a key determinant of the growth rate (11). The picture that emerges from such considerations has formed the basis of a systematic theory of bacterial growth, based on empirical "growth laws", similar to the phenomenological laws of physics (11, 13). The theory provides a successful framework for the analysis of the interdependence of cell growth and gene expression, of the effects of antibiotics, and of protein overexpression (11) without the need to characterize how the individual steps of synthesis and degradation are affected by the global state of the cell (14).

In addition to their high ribosome content, rapidly growing bacteria also contain large amounts of other proteins associated with translation. In particular, the most abundant protein in fast-growing *E. coli* is elongation factor Tu (EF-Tu), with about six to seven molecules per ribosome (15), and accounting for up to 10% of the total protein mass under conditions of rapid growth (2, 16). EF-Tu escorts charged tRNA to the ribosome and protects it from losing its amino-acylation (17, 18). The concentration of EF-Tu is closely linked to the concentration of tRNA, as EF-Tu and tRNA are typically present in cells as ternary complexes together with GTP (which activates EF-Tu) (19, 20). EF-Tu and the 21 different tRNA synthetases (21), which charge

tRNAs with the corresponding amino acids, amount to a total tRNA-affiliated protein mass of about two-thirds that of the ribosomal protein mass in moderate-to-fast growth (15). As the fraction of ribosomes allocated to making ribosomal proteins is a central factor for cell growth, synthesizing this protein mass constitutes a massive "investment" of cellular resources by diverting ribosomes away from making ribosomal proteins and toward making tRNA-affiliated proteins.

To understand why rapidly growing cells allocate such a large fraction of their proteome to tRNA-affiliated proteins, we first estimate the concentration of these proteins needed to saturate the ribosome with charged tRNA. We find that slow diffusion of the ternary complexes defines a lower bound on this concentration, pointing to the role molecular crowding plays in proteome allocation, which ultimately sets a speed limit on protein synthesis and cell growth.

The high abundance of tRNA-affiliated proteins necessary to maintain rapid translation must be balanced against the burden associated with devoting precious ribosomal resources to their synthesis (22). To elucidate how rapid growth can be achieved subject to these constraints, we develop an explicit theory that accounts for the cost of ribosomal protein synthesis and the cost of synthesizing tRNA-affiliated proteins. Our theory is based on the quantitative phenomenological framework developed by Scott et al. (11), but in contrast to the earlier work allows the speed of translation to vary (through the abundance of tRNAs and tRNA-affiliated proteins). The resulting analysis shows that this theory is consistent with a large amount of experimental data and indicates that coregulation of tRNA- and ribosome-affiliated proteins results in near-optimal resource allocation irrespective of growth rate. Furthermore, our theory provides a resolution to a long-standing contradiction in bacterial physiology, between the observed growth-rate dependence of the translation speed (23–25) (Fig. 1B) and the linear relation between ribosome concentration and growth rate traditionally attributed to a constant translation speed (1).

Significance

In rapidly growing bacteria, one-third of the proteins are ribosomal proteins and another one-sixth are ternary complexes (TCs) that bring amino acids to the ribosomes, making TCs the most abundant protein species. This study reveals that this very high TC content is forced upon the cell due to the slow diffusion of the large TCs in the crowded cytoplasm. A model is presented to describe how a bacterial cell solves the problem of proteomic resource allocation, balancing the two requirements for many ribosomes and for rapid translation. The model shows that proportional or approximately proportional synthesis of ribosomal proteins and TCs, as observed in *Escherichia coli* cells, is a near-optimal allocation scheme.

Author contributions: S.K., M.S., S.P., and T.H. designed research; S.K. and M.S. performed research; S.K., M.S., and S.P. analyzed data; and S.K., M.S., and T.H. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: klumpp@mpikg.mpg.de.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1310377110/-DCSupplemental.




Fig. 1. Translation and cell growth. (A and B) Growth-rate dependence of (A) the ribosome concentration (measured by the fraction of total protein that is ribosomal protein) and (B) the translation elongation speed for growth in media with different nutrient content. The data points in A are from refs. 2, 10, and 11, and those in B are from refs. 2, 10, and 23–25. The black lines indicate a linear and Michaelis–Menten-type fit to the data, respectively. (C) Michaelis–Menten description of the translation speed: Peptide chain elongation is described as reversible binding of a ternary complex (T) to a ribosome (R) containing a peptide chain with n amino acids and chain-irreversible chain elongation. Binding of ternary complexes to the ribosome is characterized by the Michaelis constant K_M . A lower limit for that constant is given by diffusion-limited binding of ternary complexes.

Results and Discussion

Diffusion of Ternary Complexes Limits the Speed of Translation. Peptide chain elongation is a complex process consisting of many sub-steps (26). For our purpose, however, it suffices to consider a simplified description by Michaelis–Menten kinetics (22), with reversible binding of a ternary complex to the ribosome followed by irreversible incorporation of the amino acid into the elongating chain (Fig. 1C). In particular, we do not consider the dependence of the translation speed on the translated codon (27), so the corresponding rates should be considered as averages over the translated sequences. In this picture, binding between the ternary complex and the ribosome is characterized by the Michaelis constant $K_M = (k_{off} + k_{elong})/k_{on}$, where k_{on} and k_{off} are the binding and unbinding rates of ternary complexes to the ribosome, and k_{elong} is the effective rate of peptide elongation. This constant establishes an absolute scale for the concentration of the ternary complex necessary for efficient protein synthesis; the lower the Michaelis constant is, the less ancillary protein is required. A lower bound of this concentration scale can be estimated by the diffusion limit as

$$K_M = \frac{k_{off} + k_{elong}}{k_{on}} \geq \frac{k_{elong}}{k_{on}} \geq \frac{k_{elong}}{k_{on}^{diff}}, \quad [1]$$

where we have replaced the binding rate by its upper limit, the diffusion-limited binding rate k_{on}^{diff} . The binding rate can be estimated as $k_{on}^{diff} = 4\pi Da$ for molecules of linear dimension a . For the ternary complex, $a \approx 2$ nm (20). Measured diffusion coefficients of proteins in the crowded cytoplasmic environment are in the range of 0.4 – $8 \mu\text{m}^2 \cdot \text{s}^{-1}$ (28–30). From the size dependence of the diffusion coefficient (30), we estimate the diffusion coefficient for the large ternary complex to be $D \approx 1 \mu\text{m}^2 \cdot \text{s}^{-1}$; thus $k_{on}^{diff} \sim 10 \text{ s}^{-1} \cdot \mu\text{M}^{-1}$. The elongation rate k_{elong} must exceed the largest translation speeds observed in vivo, 20–23 aa/s (2, 10). (The maximal elongation rate required here is an average over the translated sequences. Specific codons may have higher elongation rates.) Assuming $k_{elong} \approx 30 \text{ s}^{-1}$, our estimate leads to a Michaelis constant in the micromolar range ($K_M \approx 3 \mu\text{M}$),

comparable to the very high concentrations (~ 3 – $30 \mu\text{M}$) of the major tRNA species found in rapidly growing *E. coli* (31).

This concentration scale can be converted to a proteome fraction (φ_M) of tRNA-affiliated proteins (referred to henceforth as T-proteins). This class of proteins includes EF-Tu and the tRNA synthetases, which are found at constant ratio to EF-Tu in *E. coli* (15). Considering the protein mass of EF-Tu and tRNA synthetases, we obtain a scale of $\varphi_M \approx 0.02$. This is consistent with the measured proteome fraction of T-proteins in the cell, which ranges between 5% and 13% depending on growth conditions and corresponds to $\sim 60\%$ of the ribosomal protein mass (Fig. 1A). As the estimated Michaelis constant corresponds to the concentration at which translation would proceed at half of its maximal speed, these numbers indicate that the translation speed ranges between 70% and 86% of its maximum, consistent with earlier estimates for *Salmonella* (16). Our estimate of the Michaelis constant thus suggests that translation indeed operates close to the diffusion limit and that macromolecular crowding imposes a substantial cost to the cell by setting a large scale for the required allocation of T-proteins.

Growth Theory with a Four-Component Model of the Proteome. Our estimates above indicate that T-proteins may account for a substantial fraction of the total proteome. Thus, a corresponding fraction of ribosomes needs to be allocated to their synthesis. As ribosomes are critical factors for cell growth, this allocation represents a substantial investment of resources to the cell. We next develop a theory of bacterial growth that accounts for the trade-off between the costs of synthesizing ribosome- and tRNA-affiliated proteins. Our theory describes the proteome as consisting of four sectors and considers the constraints on allocating resources of protein synthesis (in particular, ribosomes) to these sectors. The theory extends the growth theory of Scott et al. (11), which was based on a three-component model of the proteome (Fig. 2A), by splitting the ribosome-affiliated proteome sector into a ribosomal fraction and a fraction of T-proteins (Fig. 2B) and allowing for a growth-rate dependence of the translation speed, which depends on the T-proteins.

We start developing our theory from the known fundamental constraints operating on exponentially growing bacteria. The first is a balance of exponential protein mass accumulation (dM/dt) by the translational output of the ribosomes,

$$dM/dt = \lambda M = k(N_{Rb} - N_{Rb,0}), \quad [2]$$

where the first equality expresses exponential growth of the cell and all its content at rate λ , and the second equality expresses the synthesis of protein by ribosomes whose total abundance is N_{Rb} (with $N_{Rb,0}$ not translating), and with the active ones working at an elongation speed k . It is convenient to express the number of ribosomes in units of the ribosomal protein mass, m_{Rb} . Then, dividing through by the total protein mass M , we arrive at a relation between the growth rate λ and the ribosomal protein fraction,

$$\lambda = \gamma \cdot (\phi_{Rb} - \phi_{Rb,0}), \quad [3]$$

where $\phi_{Rb} = N_{Rb} \cdot m_{Rb} / M$ is the total ribosomal protein mass fraction, $\phi_{Rb,0} = N_{Rb,0} \cdot m_{Rb} / M$ is the mass fraction of inactive ribosomes, and $\gamma = k / m_{Rb}$ is the rescaled elongation rate, which has been taken as constant with $k = k_{\text{elong}}$ in previous work (11, 13).

A second constraint is the need to balance the flux of energy and amino acids consumed in translation against the influx mediated by metabolic proteins (denoted by the mass fraction ϕ_P) (11),

$$\lambda = \nu \phi_P, \quad [4]$$

where ν is a phenomenological parameter that characterizes the quality of the growth medium (11).

The significant fraction of the proteome devoted to T-proteins (ϕ_T) necessary to maintain rapid translation also needs to be accounted for in the growth theory. Previously, Scott et al. used the observed approximate proportionality between ϕ_T and the ribosomal fraction; i.e.,

$$\phi_T = \alpha \cdot \phi_{Rb}, \quad [5]$$

with the observed value $\alpha \approx 0.6$ for a broad range of growth rates (discussed below), to define a ribosome-affiliated proteome fraction $\phi_R \equiv \phi_{Rb} + \phi_T = (1 + \alpha) \cdot \phi_{Rb}$. This amounts to increasing the unit mass of ribosomal proteins to $(1 + \alpha) m_{Rb}$, leading to the relation

$$\lambda = \gamma_R (\phi_R - \phi_{R,0}) \quad [6]$$

replacing Eq. 3, with $\gamma_R = k / [(1 + \alpha) \cdot m_{Rb}]$.

The dependence of the growth rate on the nutrient quality (ν) and the elongation rate (γ_R) is derived in Scott et al.'s theory by simultaneously solving Eqs. 4 and 6 subject to the constraint $\phi_R + \phi_P + \phi_Q = 1$, where ϕ_Q is a growth-rate-independent proteome fraction. (Data in ref. 11 suggest a value $\phi_Q \approx 55\%$.) The results are summarized in Fig. 2A and in *SI Text*.

In this work, we examine in detail the effect of the cell's allocation of the tRNA-affiliated fraction (T-fraction), ϕ_T , on the growth rate. To investigate the trade-off between a high translation speed and a high proteome fraction of T-proteins, the translation speed γ is now taken to depend on ϕ_T . This dependence is again described by a Michaelis–Menten relation (22),

$$\gamma(\phi_T) = \gamma_{\text{max}} \frac{\phi_T}{\phi_T + \varphi_M}. \quad [7]$$

This relation follows from the Michaelis–Menten relation of the translation speed on the ternary complex concentration c_T (22), $k = k_{\text{elong}} c_T / (K_M + c_T)$, by defining a rescaled maximal translation speed $\gamma_{\text{max}} = k_{\text{elong}} / m_{Rb}$ and by expressing concentrations in terms of the corresponding proteome fractions ($\phi_T = \sigma c_T$, with $\sigma \approx 2.5 \times 10^{-4} \mu\text{M}^{-1}$) (Methods). The Michaelis constant φ_M in Eq. 7 is thus also expressed in units of a proteome fraction; it is related to the usual Michaelis constant, K_M (estimated above), via $\varphi_M = \sigma K_M \times n_t$, where $n_t \approx 30$ is the number of different tRNAs (not counting some rare tRNAs) (Methods), such that each codon sees a tRNA concentration of K_M . The translation speed actually depends on the fraction of charged T-proteins and not the total pool ϕ_T . In moderate-to-fast growth, the charged fraction of tRNA is 70–80% (32), leading to an effective rescaling of the Michaelis constant φ_M . Here, we do not include an explicit partitioning between charged and uncharged T-proteins.

The four-component proteome model is thus defined by Eqs. 3 and 4, with γ given by Eq. 7, subject to the proteome constraint

$$\phi_{Rb} + \phi_T + \phi_P + \phi_Q = 1. \quad [8]$$

This model provides the growth rate λ and the proteome fractions ϕ_{Rb} and ϕ_P for different choices of the T-fraction ϕ_T , once the parameters ν , γ_{max} , and φ_M are specified. In the following, we assume that $\phi_T = \alpha \phi_{Rb}$, with a constant (growth-rate-independent) α , an approximation consistent with observed ratios of EF-Tu to ribosomes (see below). The (approximate) proportionality reflects the coregulation of ribosomal proteins and elongation factors, which are found in the same operons (likewise, many tRNA genes are in ribosomal RNA operons) (1).

The Four-Component Model Is Consistent with the Growth-Rate Dependence of Translation-Related Parameters.

Next, we compare with experimental data the growth-rate dependencies of the proteome fractions and of the translation speed obtained from the four-component model. This comparison also allows us to estimate the parameters φ_M and γ_{max} . We first consider the effect of changing growth by changing the nutrient quality (changing ν). By inverting the dependence of the growth rate on ν , we can plot proteome fractions and the translation speed as a function of the growth rate (rather than as a function of ν itself). As in the three-




Fig. 2. Proteome partitioning models with three and four proteome fractions. (A) Three-component model of the proteome of refs. 11 and 13: The proteome consists of three sectors, a fixed fraction (Q), a ribosome-affiliated fraction (R), which exhibits the linear growth rate dependencies for nutrient variation and translation inhibition sketched on the left, and a fraction of metabolic and other fractions (P) that has the opposite growth rate dependence. (B) The four-component model of this study splits the ribosome-affiliated fraction into a ribosomal protein fraction (Rb) and a fraction of tRNA- or translation speed-affiliated proteins (T), which increase and decrease together.

component model, there is a maximal growth rate (obtained in the limit of $\nu \rightarrow \infty$), at which the ribosomal proteome fraction is maximal [$\phi_{Rb} \rightarrow (1 - \phi_Q)/(1 + \alpha)$] and the metabolic proteome fraction vanishes ($\phi_P \rightarrow 0$, a growth-rate-independent basal offset in the metabolic sector that accounts for the metabolic activity still present at maximal growth can be absorbed into the growth-rate-independent Q sector; for a discussion see the supporting information of ref. 33). For smaller growth rates, ϕ_{Rb} depends approximately linearly on the growth rate λ ,

$$\phi_{Rb} = \phi_{Rb,0} + \frac{\varphi_M}{\alpha} + \frac{\lambda}{\gamma_{\max}} (1 - f(\lambda)), \quad [9]$$

up to a small correction $f(\lambda)$ (Eq. S10), which turns out to be negligible for moderate to fast growth (e.g., above $\sim 0.5/h$). However, comparing Eq. 9 to the form [6] for the three-component model, the interpretations of the slope and the offset of the linear relation are different as is discussed below. Because $\phi_T \propto \phi_{Rb}$, ϕ_T also depends linearly on λ , and consequently the translation speed γ (Eq. 7) is approximately given by a Michaelis-Menten function of growth rate for moderate-to-fast growth.

In Fig. 3 A and B, we plot the calculated growth rate dependence of the ribosomal proteome fraction (ϕ_{Rb}) and the translation speed $k = \gamma(\phi_T)/m_{Rb}$, matched to the corresponding data for growth in different nutrients (data as in Fig. 1 A and B). With α set to 0.6, this matching determines the two parameters characterizing the translation speed, the maximal speed γ_{\max} and the Michaelis constant φ_M , and the offset $\phi_{Rb,0}$ attributed to inactive ribosomes. The values of the fitted parameters are given in Table S1. Specifically, we obtain $\varphi_M = 0.029$, which is only slightly larger than the minimal value (0.02) estimated above based on diffusion-limited ternary complex binding. As a consistency check we also compare the calculated growth rate dependence of the T-fraction (ϕ_T) with an estimate of that proteome fraction from the measured amount of EF-Tu for growth in different nutrients (15, 34) (the measured proteome fraction of EF-Tu was multiplied by 1.6 to account for the tRNA synthetases) (Methods). This estimate and the results from the model show good agreement, with exception of the fastest growth.

Finally, we plotted the ratio $\phi_T : \phi_R$ in Fig. 3D, expressed as a mass ratio (α) and as a ratio of molecule numbers (EF-Tu per ribosome) together with experimental data for EF-Tu per ribosome (15, 34, 35) (gray symbols) and tRNA per ribosome (31, 36) (white symbols). Within the present model the ratio $\phi_T : \phi_R$ is constant; $\alpha = 0.6$ corresponds to 7.6 elongation factors per ribosome. The data for EF-Tu per ribosome show a slight, but systematic reduction with increasing growth rate, possibly indicating that the two proteome fractions are not exactly coregulated, at least during very fast growth. For the ratio of tRNAs to ribosomes, the situation is less clear: There is a slight decrease at fast growth in one dataset (31), but not in another (1, 36). The increased tRNA content seen at very slow growth [based on data from chemostat growth (36)] is believed to arise from differential stability of tRNA and rRNA at slow growth (2) and thus may not reflect the dependence of the corresponding proteome fraction. In sum, over moderate-to-fast growth rates, the four-component model with coregulation of T-proteins and ribosomal proteins recapitulates the growth dependence of the proteins involved in translation, as well as the growth-rate-dependent translation speed under conditions where growth is modulated by nutrient quality.

We next calculated the growth-rate dependence of the ribosomal and T-fractions under translation-limited growth, as studied in ref. 11 using ribosome-targeting antibiotics, which is implemented in the model by reducing the maximal translation speed γ_{\max} for constant ν (dashed red lines in Fig. 3). With decreasing γ_{\max} , the ribosomal fraction increases as observed with increasing antibiotic concentrations (11). It exhibits a linear dependence on the growth rate and extrapolates to a maximal ribosomal fraction that is independent of the nutrient quality ν , i.e., independent





Fig. 3. Growth-rate dependence of translation-related parameters. (A) Ribosomal proteome fraction; (B) translation speed; (C) tRNA-affiliated proteome fraction; (D) ratio of tRNA-affiliated and ribosomal fractions (as mass ratio α and ratio of numbers of molecules). In all plots, the solid line is for growth variation by the nutrient quality (varying ν), and the dashed lines are for growth variation by translational inhibition, e.g., using antibiotics (varying γ_{\max}). The parameters (Table S1) are chosen to match the experimental data (gray symbols). Data sources: (A and B) As in Fig. 1 C and D; T-protein data are from refs. 15, 34, and 35; tRNA data in D are from refs. 31 and 36.

of the growth medium in the absence of the antibiotic (Fig. 3A). The tRNA-affiliated fraction ϕ_T also increases, being forced to do so by the assumption of coregulation (Fig. 3C). We note that in the four-component model, translation limitation can also be obtained by increasing the Michaelis constant of translation φ_M , which leads to the same growth-rate dependence as changing the maximal translation speed γ_{\max} .

Coregulation of Ribosomal and tRNA-Affiliated Proteins Corresponds to Near-Optimal Allocation of Protein Synthesis Resources. So far, we have assumed that the factor α , the ratio between the tRNA-affiliated and ribosomal proteome fractions, is constant and independent of the growth rate. Based on an optimization scheme originally proposed by Ehrenberg and Kurland (22), we have varied α and adjusted it such that the growth rate is maximized for any given growth conditions, i.e., for fixed medium parameters (ν , γ_{\max} , φ_M). The optimal α is dependent on the growth rate and a systematic decrease of the ratio of T-proteins to ribosomes with the growth rate is obtained (Fig. 4). Although the data for EF-Tu per ribosome (symbols in Fig. 4C) show a slight decrease with increasing growth rate, the optimal α is smaller than the observed values. However, for all realistic values of the growth rate (up to three doublings per hour), a constant $\alpha = 0.6$ as used above leads to a growth rate within 10% of the maximum obtained by optimizing α , and the experimental ratio of EF-Tu per ribosome, which decreases slightly at fast growth, even remains in the $\pm 5\%$ region (gray symbols in Fig. 4C). As a result, we conclude that coregulation (or approximate coregulation) of ribosomal and T-proteins is a near-optimal strategy for the cell, which may be “good enough” to achieve fast growth. As the additional complexity of regulating the two proteome sectors separately to achieve full optimization may incur additional fitness costs, coregulation (with possibly some adjustment to lower the ratio ϕ_T/ϕ_R at fast growth) may even be the most robust strategy for the cell to adopt.

Effect of the Michaelis Constant on Cell Growth. Finally, we turn to the effective Michaelis constant (φ_M or K_M) for ternary complex



binding to the ribosome, which plays a unique role in our model: It sets an absolute concentration scale for the T-fraction, whereas otherwise the model is concerned with allocating fractions of the total proteome. Fig. 5A shows the maximal growth rate as a function of the Michaelis constant, with a fixed value of α and with α optimized for each given value of φ_M (red and blue lines, respectively). For fixed α , lowering the Michaelis constant increases the translation speed, and thus the growth rate, by saturating ribosomes with ternary complexes. Optimizing α with respect to a smaller Michaelis constant allows the cell to reduce its investment in the T-fraction and to allocate more resources to the ribosomal fraction (Fig. 5B), as with a smaller φ_M the same translation speed can be achieved with a smaller concentration of ternary complexes. In either case, a decrease of the Michaelis constant increases the growth rate. Above we have argued that the value of the Michaelis constant is set by a physical limit, namely that binding of ternary complexes to the ribosome cannot be faster than diffusion-limited. This limit is indicated in Fig. 5 by the dashed vertical line. It points toward a key role molecular crowding plays in cell growth. The link between crowding and cell growth may be complex (37), as increased crowding slows diffusion (38), but also reduces unbinding (39) and effectively increases concentrations by reducing the accessible volume. However, if translation remains close to the diffusion limit, as we suggest here, slow diffusion should be the dominant effect of crowding (40, 41).

Mechanistic Interpretation of the Phenomenological Model. We have noted above that our growth theory with a four-component model of the proteome leads to a modified interpretation of the slope ($1/\gamma$) of the linear relation between ribosome concentration and growth rate (Fig. 1A). We now discuss this point in more detail and show that it resolves an old controversy surrounding the phenomenological approach to bacterial growth. Within the phenomenological approach, the mass balance argument presented in Eqs. 2 and 3 was used to identify the phenomenological parameter

γ with a growth-rate-independent translation speed (1). This interpretation is consistent with the observation that the slope is changed in mutants with slow ribosomes, with a strong correlation between the inverse slopes and the corresponding translation speeds measured in vitro (11). Nevertheless, that interpretation stood in contradiction to direct measurements in *E. coli* that indicate a dependence of the translation speed on growth rate (23–25) (Fig. 1B). Our discussion of the four-component model above (Eq. 9) shows that the growth law of Fig. 1A is consistent with a growth-rate-dependent translation speed, with the modified interpretation that the slope in Fig. 1A be identified with a growth-rate-independent maximal translation speed, γ_{\max} (Eq. 6). The modified interpretation remains consistent with the slope-speed correlation for the slow ribosome mutants. In fact, the need for such modified interpretation is also obtained by asking for a functional form of $\gamma(\lambda)$ that is compatible with the linear relation between ribosome concentration and growth rate. This argument, which is made in *SI Text*, leads to a Michaelis-Menten-type dependence of the translation speed on growth rate (Eq. 9 without the nonlinear term, included as the black line in Fig. 1B with parameters given in Table S1).

Quantitative phenomenology provides a powerful approach for predicting physiological responses to a variety of perturbations (11) and for uncovering regulatory links within their physiological context (33). However, our present study indicates that care must be taken when assigning a mechanistic interpretation to the empirical parameters appearing in such models. Coarse graining can bring into focus relations that are not apparent in rich datasets; nevertheless, coarse graining is a projection of the underlying dynamics of the system, and different (and, possibly, mutually contradictory) mechanistic models may be consistent with a particular set of phenomenological relations.

Concluding Remarks. In this study, we have derived an explicit model for proteome partitioning that accounts for the fact that an increase of the translation speed incurs a cost to the cell in the form of required protein such as elongation factors and tRNA synthetases. This fraction of the total protein mass that is devoted to the maintenance of the translation speed is ultimately governed by the slow diffusion of the large ternary complexes in a crowded cytoplasm. Furthermore, the analysis resolves the apparent discrepancy between a very successful phenomenological framework for understanding the interdependence of cell growth and gene expression based on the linear relation between ribosome concentration and growth rate (which is naturally interpreted as reflecting a constant translation speed) on the one hand and the measured growth rate dependence of the translation speed on the other hand. We have shown that a growth-rate dependence of the translation speed is consistent with the phenomenological approach and results in only a modified interpretation of the parameters.

The limitation of the Michaelis constant for ternary complex-ribosome binding by slow diffusion points toward a key role for molecular crowding in proteomic resource allocation and cell growth. In bacterial cells, the degree of crowding (or the water content of a cell) can be varied via the osmolarity of the growth




Fig. 5. (A and B) Diffusion limitation of the minimal Michaelis constant. (A) Maximal growth rate and (B) ribosomal and tRNA-affiliated proteome fractions ϕ_{Rb} and ϕ_T (solid and dashed lines, respectively) as a function of the Michaelis constant (φ_M or K_M) of binding of ternary complexes to ribosomes. The red lines are for constant α and the blue lines for optimal α such that the growth rate is maximized for each value of the Michaelis constant. The dashed lines indicate our estimate of the diffusion-limited minimal value of the Michaelis constant.

medium. Cells adapted to reduced water content still exhibit exponential growth, but with a reduced growth rate (42). Studying the ribosome content and translation speed of these cells should provide valuable insight into the link between crowding and cell growth. Specifically, our study suggests that the slow diffusion of ternary complexes is an important contribution to the slowdown of growth. This could be directly probed by following the diffusion of ternary complexes in cells under osmotic challenge.

Methods

Growth Theory with a Four-Component Proteome Partitioning Model. The growth theory with the four-component model of the proteome is defined by Eqs. 3–5, 7, and 8. Solving these equations defines the growth rate (λ), the proteome fractions (ϕ_{Rb} , ϕ_T , ϕ_P), and the translation speed (γ) as functions of the parameters of translation (γ_{max} , ϕ_M) and the parameter ν , which characterizes the nutrient quality of the medium. Inverting the dependence of λ on ν or γ_{max} leads to the growth-rate dependence of the quantities for nutrient-limited and translation-limited growth, respectively. A mathematical summary of the theory with a three-component and a four-component model of the proteome is given in *SI Text*.

Conversion of Concentrations and Proteome Fractions. Proteome fractions (ϕ) and concentrations (c) are converted through $\phi = \sigma c$. The factor σ is obtained as

$$\sigma = \sigma_0 \frac{\text{volume}}{\text{mass}} \frac{\text{mass}}{\text{total protein}} N_{aa} \approx 3.8 \times 10^{-7} \mu\text{M}^{-1} \times N_{aa}, \quad [10]$$

- Maaloe O (1979) Regulation of the protein-synthesizing machinery - ribosomes, tRNA, factors, and so on. *Biological Regulation and Development*, ed Goldberger RF (Plenum, New York), Vol 1, pp 487–542.
- Bremer H, Dennis PP (1996) Modulation of chemical composition and other parameters of the cell by growth rate. *Escherichia coli and Salmonella*, ed Neidhardt FC (Am Soc Microbiol, Washington, DC), 2nd Ed, pp 1553–1569.
- Russell JB, Cook GM (1995) Energetics of bacterial growth: Balance of anabolic and catabolic reactions. *Microbiol Rev* 59:48–62.
- Dennis PP, Ehrenberg M, Bremer H (2004) Control of rRNA synthesis in *Escherichia coli*: A systems biology approach. *Microbiol Mol Biol Rev* 68:639–668.
- Schaechter M, Maaloe O, Kjeldgaard NO (1958) Dependency on medium and temperature of cell size and chemical composition during balanced growth of *Salmonella typhimurium*. *J Gen Microbiol* 19:592–606.
- Warner JR (1999) The economics of ribosome biosynthesis in yeast. *Trends Biochem Sci* 24:437–440.
- Vind J, Sorensen MA, Rasmussen MD, Pedersen S (1993) Synthesis of proteins in *Escherichia coli* is limited by the concentration of free ribosomes. Expression from reporter genes does not always reflect functional mRNA levels. *J Mol Biol* 231:678–688.
- Stoebel DM, Dean AM, Dykhuizen DE (2008) The cost of expression of *Escherichia coli* lac operon proteins is in the process, not in the products. *Genetics* 178:1653–1660.
- Neidhardt FC, Magasanik B (1960) Studies on the role of ribonucleic acid in the growth of bacteria. *Biochim Biophys Acta* 42:99–116.
- Forchhammer J, Lindahl L (1971) Growth rate of polypeptide chains as a function of the cell growth rate in a mutant of *Escherichia coli* 15. *J Mol Biol* 55:563–568.
- Scott M, Gunderson CW, Mateescu EM, Zhang X, Hwa T (2010) Interdependence of cell growth and gene expression: Origins and consequences. *Science* 330:1099–1102.
- Maaloe O (1969) An analysis of bacterial growth. *Dev Biol* 3(Suppl):33–58.
- Scott M, Hwa T (2011) Bacterial growth laws and their applications. *Curr Opin Biotechnol* 22:559–565.
- Klump S, Zhang Z, Hwa T (2009) Growth rate-dependent global effects on gene expression in bacteria. *Cell* 139:1366–1375.
- Neidhardt FC, Bloch PL, Pedersen S, Reeh S (1977) Chemical measurement of steady-state levels of ten aminoacyl-transfer ribonucleic acid synthetases in *Escherichia coli*. *J Bacteriol* 129:378–387.
- Tubulekas I, Hughes D (1993) Growth and translation elongation rate are sensitive to the concentration of EF-Tu. *Mol Microbiol* 8:761–770.
- Nierhaus KH (2004) The elongation cycle. *Protein Synthesis and Ribosome Structure*, eds Nierhaus KH, Wilson DN (Wiley-VCH, Weinheim, Germany), pp 323–366.
- Andersen C, Wiborg O (1994) *Escherichia coli* elongation-factor-Tu mutants with decreased affinity for aminoacyl-tRNA. *Eur J Biochem* 220:739–744.
- Gouy M, Grantham R (1980) Polypeptide elongation and tRNA cycling in *Escherichia coli*: A dynamic approach. *FEBS Lett* 115(2):151–155.
- Krab IM, Parmeggiani A (1998) EF-Tu, a GTPase odyssey. *Biochim Biophys Acta* 1443(1–2): 1–22.
- Martinis SA, Schimmel P (1996) Aminoacyl-tRNA synthetases: General features and relationships. *Escherichia coli and Salmonella*, ed Neidhardt FC (Am Soc Microbiol, Washington, DC), pp 887–901.
- Ehrenberg M, Kurland CG (1984) Costs of accuracy determined by a maximal growth rate constraint. *Q Rev Biophys* 17(1):45–82.
- Dalbow DG, Young R (1975) Synthesis time of beta-galactosidase in *Escherichia coli* B/r as a function of growth rate. *Biochem J* 150(1):13–20.
- Young R, Bremer H (1976) Polypeptide-chain-elongation rate in *Escherichia coli* B/r as a function of growth rate. *Biochem J* 160(2):185–194.
- Pedersen S (1984) *Escherichia coli* ribosomes translate in vivo with variable rate. *EMBO J* 3:2895–2898.
- Wintermeyer W, et al. (2004) Mechanisms of elongation on the ribosome: Dynamics of a macromolecular machine. *Biochem Soc Trans* 32:733–737.
- Klump S, Dong J, Hwa T (2012) On ribosome load, codon bias and protein abundance. *PLoS ONE* 7:e48542.
- Elowitz MB, Surette MG, Wolf PE, Stock JB, Leibler S (1999) Protein mobility in the cytoplasm of *Escherichia coli*. *J Bacteriol* 181(1):197–203.
- Elf J, Li GW, Xie XS (2007) Probing transcription factor dynamics at the single-molecule level in a living cell. *Science* 316:1191–1194.
- Kumar M, Mommer MS, Sourjik V (2010) Mobility of cytoplasmic, membrane, and DNA-binding proteins in *Escherichia coli*. *Biophys J* 98:552–559.
- Dong H, Nilsson L, Kurland CG (1996) Co-variation of tRNA abundance and codon usage in *Escherichia coli* at different growth rates. *J Mol Biol* 260:649–663.
- Sorensen MA (2001) Charging levels of four tRNA species in *Escherichia coli* Rel(+) and Rel(-) strains during amino acid starvation: A simple model for the effect of ppGpp on translational accuracy. *J Mol Biol* 307:785–798.
- You C, et al. (2013) Coordination of bacterial proteome with metabolism by cyclic AMP signalling. *Nature* 500:301–306.
- Pedersen S, Bloch PL, Reeh S, Neidhardt FC (1978) Patterns of protein synthesis in *Escherichia coli*. Catalog of amount of 140 individual proteins at different growth rates. *Cell* 14(1):179–190.
- Reeh S, Pedersen S (1978) Regulation of *Escherichia coli* elongation factor synthesis in vivo. *Gene Expression (11th FEBS Meeting Copenhagen 1977)*, eds Clark BFC, Klenow H, Zeuthen J (Pergamon, Oxford), pp 89–98.
- Jacobsen H (1974) Peptide chain growth rate, cell size, protein, DNA, tRNA and rRNA as function of the growth rate in chemostat and batch cultures of *E. coli*. PhD thesis (University of Copenhagen, Copenhagen).
- Tadmor AD, Tlusty T (2008) A coarse-grained biophysical model of *E. coli* and its application to perturbation of the rRNA operon copy number. *PLoS Comput Biol* 4: e1000038.
- Konopka MC, et al. (2009) Cytoplasmic protein mobility in osmotically stressed *Escherichia coli*. *J Bacteriol* 191:231–237.
- Morelli MJ, Allen RJ, Wolde PR (2011) Effects of macromolecular crowding on genetic networks. *Biophys J* 101:2882–2891.
- Zimmerman SB, Minton AP (1993) Macromolecular crowding - biochemical, biophysical, and physiological consequences. *Annu Rev Biophys Biomol Struct* 22:27–65.
- Ellis RJ (2001) Macromolecular crowding: Obvious but underappreciated. *Trends Biochem Sci* 26:597–604.
- Cayley S, Record MT (2004) Large changes in cytoplasmic biopolymer concentration with osmolality indicate that macromolecular crowding may regulate protein-DNA interactions and growth rate in osmotically stressed *Escherichia coli* K-12. *J Mol Recognit* 17:488–496.
- Klump S (2011) Growth-rate dependence reveals design principles of plasmid copy number control. *PLoS ONE* 6:e20403.
- Nanninga N, Woldringh CL (1985) Cell growth, genome duplication, and cell division. *Molecular Cytology of Escherichia coli*, ed Nanninga N (Academic, London), pp 259–318.

Supporting Information

Klumpp et al. 10.1073/pnas.1310377110

SI Text

Summary of Growth Theory with Three- and Four-Component Proteome Models

Growth Theory with a Three-Component Model of the Proteome. We briefly summarize the growth theory of Scott et al. (1) that is based on a three-component model of the proteome. This model partitions the total proteome of a cell into three sectors, R, P, and Q that represent ribosomal and ribosome-affiliated proteins, metabolic proteins, and a fixed fraction of proteins that are expressed in a growth-rate-independent fashion, respectively (Fig. 2A). The mass fractions of the total protein mass that these three sectors account for (denoted by ϕ_R , ϕ_P , ϕ_Q , respectively) have to add up to one,

$$\phi_R + \phi_P + \phi_Q = 1. \quad [\text{S1}]$$

The three sectors exhibit different growth rate dependencies: The Q sector is independent of growth rate, includes proteins regulated by negative feedback (2), and represents about 55% of the protein mass. The ribosomal sector increases linearly with growth rate if growth is modulated by changing the nutrient quality,

$$\phi_R = \phi_{R,0} + \frac{\lambda}{\gamma_R}, \quad [\text{S2}]$$

where γ_R is the speed of translation (Eq. 6 in the main text). The growth-rate-independent offset $\phi_{R,0}$ can be interpreted as representing inactive ribosomes. The metabolic sector containing proteins for nutrient uptake and processing increases linearly with growth rate under conditions of translational inhibition,

$$\phi_P = \frac{\lambda}{\nu}. \quad [\text{S3}]$$

The two empirical relations given by Eqs. S2 and S3 can be rationalized as representing a balance of resource allocation to either protein synthesis or nutrient uptake. To achieve exponential growth, both relations [S2] and [S3] must be satisfied simultaneously. As a consequence, the growth rate can be considered as a function of the medium parameters γ_R and ν , which is fully determined by Eqs. S1–S3 and given by a Michaelis–Menten-type relation,

$$\lambda(\gamma_R, \nu, \phi_Q) = (1 - \phi_Q) \frac{\nu \gamma_R}{(\nu + \gamma_R)}. \quad [\text{S4}]$$

In addition to the parameters γ_R and ν , the growth rate depends upon the size of the Q sector, which could also be a medium-dependent parameter, but for moderate-to-fast growth of *Escherichia coli*, it is found to be approximately independent of the medium (1).

Growth Theory with a Four-Component Model of the Proteome and Coregulation of tRNA- and Ribosome-Affiliated Proteins. To allow for a growth-rate-dependent speed of translation we extend the three-component model by introducing a fourth component of the proteome that contains tRNA-affiliated proteins (T-proteins), in particular elongation factor Tu, the most abundant protein in the cell. The R-protein sector in the three-component model is then partitioned between ribosomal proteins, with mass fraction ϕ_{Rb} , and tRNA-affiliated proteins, with mass fraction ϕ_T , so that

$\phi_R = \phi_{Rb} + \phi_T$. The relation between the mass fraction of ribosomal proteins and the growth rate is modified by introducing a translation speed that exhibits a Michaelis–Menten dependence on the size of the T-fraction ϕ_T (Eq. 7 in the main text),

$$\gamma(\lambda) = \frac{\gamma_{\max} \phi_T}{\phi_T + \varphi_M}, \quad [\text{S5}]$$

such that

$$\lambda = (\phi_{Rb} - \phi_{Rb,0}) \frac{\gamma_{\max} \phi_T}{\phi_T + \varphi_M}. \quad [\text{S6}]$$

Here φ_M is the Michaelis–Menten constant expressed in units of proteome fraction. It is related to the usual Michaelis–Menten constant (in units of concentration), K_M , via $\varphi_M = \sigma K_M \times n_t$, where $n_t \approx 30$ is the effective number of different tRNAs (not counting the rarest tRNAs), such that each codon sees a tRNA concentration of K_M . The factor σ is the same as in Eq. 10 in the main text.

With a fourth protein sector, the partitioning constraint becomes

$$\phi_P + \phi_Q + \phi_{Rb} + \phi_T = 1. \quad [\text{S7}]$$

To fully determine the dependence of the growth rate on the medium parameters we need one more equation, which is obtained from the assumption of coregulation,

$$\phi_T = \alpha \phi_{Rb}, \quad [\text{S8}]$$

where α is a constant, obtained from the mass ratio of proteins known to belong to the T-fraction (elongation factors Tu and Ts, tRNA synthetases) and the ribosome-affiliated fraction [ribosomal proteins, elongation factor (EF)-G]. Counting elongation factors Tu and Ts as well as tRNA synthetases in the T sector and ribosomal proteins and elongation factor G in the ribosomal sector, we can estimate $\alpha \approx 0.6$. With the assumption of coregulation, the four-component model is effectively reduced to a three-component model with Eq. S1 replaced by Eq. S7, which becomes $\phi_P + \phi_Q + (1 + \alpha)\phi_{Rb} = 1$, and Eq. S2 replaced by

$$\phi_{Rb} = \phi_{Rb,0} + \frac{\varphi_M}{\alpha} + \frac{\lambda}{\gamma_{\max}} (1 - f(\lambda)), \quad [\text{S9}]$$

which is obtained by solving Eq. S6 for ϕ_{Rb} , using Eq. S8. In this expression, the first three terms correspond to Eq. S14 below, which is obtained with a Michaelis–Menten-type dependence of the translation speed on the growth rate (with an additional offset φ_M/α and $\lambda_M = \varphi_M \gamma_{\max}/\alpha$). The last term, with

$$f(\lambda) = \left(1 + 2\lambda_M/\lambda - \sqrt{1 + 4\lambda_M/\lambda}\right) / 2, \quad [\text{S10}]$$

is a correction most relevant at slow growth, which is small for fast growth rates (i.e., faster than 1 doubling in 85 min).

Solving Eqs. S3, S7, and S8 leads to a generalization of the Michaelis–Menten equation for growth (Eq. S4) that expresses the growth rate as function of the medium parameters, ν , γ_{\max} , φ_M , and of ϕ_Q . The maximal growth rate (for given translation characteristics, γ_{\max} , φ_M) is given by

$$\lambda_{\max} = \frac{\gamma_{\max}^2 (1 - \phi_Q)^2}{[\lambda_M (1 + \alpha) + \gamma_{\max} (1 - \phi_Q)]}. \quad [\text{S11}]$$

Combining the dependence of the growth rate on the medium parameters with Eqs. S3, S6, and S8 determines the proteome fractions as functions of the medium parameters.

Conditions for a Linear Relation Between Ribosome Concentration and Growth Rate

We asked under what conditions a growth-rate-dependent elongation speed would be consistent with the very robust linear relation observed between the ribosome concentration and the growth rate (Fig. 1A). Such consistency requires that the growth-rate-dependent translation speed, $\gamma(\lambda)$, obeys

$$\phi_{Rb}(\lambda) - \phi_{Rb,0} = \frac{\lambda}{\gamma(\lambda)} = a + b\lambda, \quad [\text{S12}]$$

where the first equality expresses the synthesis of total protein by ribosomes (Eq. 3 in the main text) and the second equality is the requirement for the observed linear dependence on growth rate. Solving this equation for the growth-rate-dependent translation speed $\gamma(\lambda)$ leads to a Michaelis–Menten-type dependence of the ribosome speed on the growth rate,

$$\gamma(\lambda) = \frac{\gamma_{\max} \lambda}{\lambda_M + \lambda}, \quad [\text{S13}]$$

where the maximal translation speed γ_{\max} and the characteristic growth rate λ_M can be related to the constants a and b in Eq. S12. Eq. S13 agrees well with the measured growth-rate dependence of the translation speed (included as the black line in Fig. 1B), with $\lambda_M = 0.44 \text{ h}^{-1}$ and $\gamma_{\max} = 10.4 \text{ h}^{-1}$, corresponding to a maximal translation speed of 25 aa/s. These parameters are similar to what we obtained by fitting the full model (Table S1). A growth-rate-dependent translation speed that follows this form leads to a linear relation between growth rate and ribosome concentration as in Eq. 3 in the main text,

$$\phi_{Rb} = \phi_{Rb,0} + \varphi_M + \frac{\lambda}{\gamma_{\max}}, \quad [\text{S14}]$$

but notably with a modified interpretation of the two parameters in that linear relation. Most importantly, the slope ($1/\gamma_{\max}$) now corresponds to the inverse maximal translation speed rather than to a constant (growth-rate-independent) translation speed. In addition, there are two contributions to the offset, one reflecting inactive ribosomes ($\phi_{Rb,0}$) and one that emerges from the growth-rate dependence of the translation speed and reflects the growth rate at which the translation speed becomes half-maximal ($\varphi_M = \lambda_M/\gamma_{\max}$).

The explicit growth-rate dependence, λ , in the translation speed, Eq. S13, can be replaced by any variable with a strong positive correlation with growth rate without abrogating the linear relation between ribosome content and growth rate. As we show in the main text, the proteome fraction of tRNA-affiliated proteins can play precisely this role.

1. Scott M, Gunderson CW, Mateescu EM, Zhang Z, Hwa T (2010) Interdependence of cell growth and gene expression: Origins and consequences. *Science* 330:1099–1102.

2. Klumpp S, Zhang Z, Hwa T (2009) Growth rate-dependent global effects on gene expression in bacteria. *Cell* 139:1366–1375.

Table S1. Parameters of the growth-rate dependence of the translation speed

	Fit to growth theory	Fit with Michaelis–Menten dependence	Comments
Maximal elongation speed, $k_{\text{elong}} \times 1 \text{ aa}$	23.5 aa/s	25 aa/s	
Maximal translation speed, γ_{\max}	10.2 h^{-1}	10.4 h^{-1}	$= k_{\text{elong}}/N_{\text{aa,ribos}}$ with $N_{\text{aa,ribos}} = 8,290$, no. amino acids per ribosome, including EF-G
Michaelis constant of translation, K_M	3.9 μM		Calculated from φ_M
Michaelis constant in proteome fraction units, φ_M	0.029		
Michaelis constant for growth rate, λ_M	0.49 h^{-1}	0.44 h^{-1}	$\lambda_M = \gamma_{\max} \varphi_M / \alpha$
Offset for ribosomal fraction, inactive ribosomes, $\phi_{Rb,0}$	0.035		
Constant proteome fraction, ϕ_Q	0.55		Including the constant offsets of the ribosomal and tRNA-related fractions, $(1 + \alpha)\phi_{Rb,0}$